

ASSESSING ENVIRONMENTAL STRESS ON THE LOGGERHEAD SEA TURTLE  
(CARETTA CARETTA) IN VIRGINIA WATERS

---

A Thesis

Presented to

The Faculty of the School of Marine Science  
The College of William and Mary in Virginia

In Partial Fulfillment  
of the Requirements for the Degree of  
Master of Arts

---

by


Sarah A. Bellmund

1988


APPROVAL SHEET


This thesis is submitted in partial fulfillment of  
the requirements for the degree of


Master of Arts


  
Sarah Anne Bellmund


Approved, December 1988

  
John A. Musick, Chairman

  
Eugene M. Burrison

  
David Zwerner

  
Robert J. Huggett

  
Robert H. George, DVM

## DEDICATION

This work is dedicated to the Watermen of the Chesapeake Bay for their knowledge, appreciation, and love of the Bay, especially, Otis Carney, Kenny Ethridge, Ed Boyd, Walter Cole Burroughs, and particularly Richard Erdt and Talmadge and Fred Jett for sharing turtles, knowledge of turtle biology, and knowledge of the Bay and especially for making this work so enjoyable.

# TABLE OF CONTENTS

	<u>Page</u>
DEDICATION . . . . .	iii
ACKNOWLEDGMENTS . . . . .	iv
LIST OF TABLES . . . . .	v
LIST OF FIGURES . . . . .	vi
ABSTRACT . . . . .	vii
INTRODUCTION . . . . .	2
METHODS . . . . .	8
RESULTS . . . . .	18
Preliminary Evaluation . . . . .	18
Multi-Variate Analysis . . . . .	40
DISCUSSION . . . . .	54
Underlying General Trends . . . . .	54
Geographic Variations . . . . .	54
Seasonal Variations . . . . .	58
Pathological Changes . . . . .	62
Multi-Variate Analyses . . . . .	63
CONCLUSIONS . . . . .	69
APPENDICES . . . . .	71
LITERATURE CITED . . . . .	73
VITA . . . . .	82



## ACKNOWLEDGMENTS

Special thanks to my thesis committee Dave Zwerner, Gene Burreson, Bob George, Bob Huggett and Jack Musick for help, advise, and support. I would like to thank those people at VIMS who gave time and support particularly, Howard Kator, Nita Walker, Lee Ellis, John Boon, Che Wu Su, Herb Austin, Bill DuPaul, Jim Owens, Paul Gerdes, Joe Desfosse, Mary Sue Mustafa, Pat Hall, Gary Anderson, Jane Lopez, Joe Choromanski, Dennis Thoney, the Ichthons from Jefferson, and Rob Hale.

I would especially like to thank David Owens and Thane Wibbles at Texas A&M University for their time to run corticosterone samples. I would also like to thank Elliot Jacobson, Bill Dunson, Frank Mazzotti, Dan Odell, and Gordon Ultsch, for review, comment and suggestion on various stages of this project. Ray Morgan and Dave Sager (Univ. MD, HPEL) supported my work and were very helpful. Peter Lutz (Univ. of Miami, RSMAS) provided time, background, and direction. Special thanks to George Barns at Walter Reed Memorial Hospital for supervision of the samples and their analysis. Special thanks also to the Gloucester Veterinary Clinic for much time and help.

I would also like to thank the many friends who went on turtle runs etc. especially turtle people past and present particularly John and Deb Keinath, Richard Byles, Molly Lutcavage, Gloria Rowe and particularly RuthEllen. I would also like to specially thank Chris Pague for much aid, support, encouragement and herp trips, Jared

Brandwien for being there when I needed him, Pat Dolan and Patrice Mason for support through hard times, Brett Burdick for songs and dead turtles, Paul DeFur for Jim Carrol, and finally the BUBBAs collectively for everything, anything or nothing, depending on case. Joel VanArman and Lynn Gulick provided invaluable support and assistance in the final preparation of this work. I would especially like to thank George Vadas, for without whom I would not have made it through.

Support was provided for this work and my time by the National Marine Fisheries Service, NE Region; the Virginia Game Commission, Non-Game Fund; and the Department of Defense, EMPRESS project.

## LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Apparent Causes of Death in Virginia Sea Turtles Examined between 1979-1983 . . . . .	4
2. Sea Turtle Morphometric Measurements . . . . .	12
3. <u>Caretta caretta</u> serum chemistries, means and standard deviations by month . . . . .	19
4. <u>Caretta caretta</u> serum chemistries, means and standard deviations by zone . . . . .	20
5. <u>Caretta caretta</u> serum chemistries, means, and standard deviations by sex . . . . .	21
6. <u>Caretta caretta</u> serum chemistries, means and standard deviations by size . . . . .	22
7. <u>Caretta caretta</u> serum chemistries, means and standard deviations by condition . . . . .	23
8. General statistics for turtles included in Q-Mode factor analysis . . . . .	43
9. Variable loadings for Factors calculated in Q-Mode factor analysis . . . . .	44
10. Endmember groups and serum profiles for Factor 1 . . . . .	48
11. Endmember groups and serum profiles for Factor 2 . . . . .	49
12. Intermediate Turtles Described by Factor Analysis . . . . .	50
13. Sera Profiles for <u>Caretta</u> with Low Body Temperatures (<18 °C) . . . . .	53

# LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Zones for Virginia sea turtles sampling . . . . .	9
2. Northern and Southern areas for blood comparisons . . . . .	10
3. Monthly mean serum protein values for all healthy turtles 1983-1985 . . . . .	24
4. Monthly mean serum electrolytes for all healthy turtles 1983-1985 . . . . .	25
5. Monthly mean serum organic components for all healthy turtles 1983-1985 . . . . .	26
6. Seasonal differences in mean values for serum proteins of <u>Caretta</u> from different regions 1983-1985 . . . . .	28
7. Seasonal differences in mean values for serum ions of <u>Caretta</u> from different regions 1983-1985 . . . . .	29
8. Seasonal differences in mean values for serum organic components of <u>Caretta</u> from different regions 1983-1985 .	30
9. Seasonal relationship between calcium and phosphorus 1983-1985 . . . . .	32
10. Seasonal relationship between sodium and albumin 1983-1985 . . . . .	34
11. Seasonal relationship between sodium and chloride 1983-1985 . . . . .	35
12. Seasonal relationship between potassium and blood urea nitrogen (BUN) . . . . .	37
13. Seasonal values for potassium . . . . .	38
14. Changes in glucose and urea nitrogen at 20°C and 26°C over time . . . . .	39
15. Mean seasonal serum corticosterone in apparently healthy <u>Caretta</u> 1984-1985 . . . . .	41
16. Varimax two factor solution for turtle serum profiles . . . .	46
17. Groupings marked on Varimax factor plot of serum profiles . .	47

## ABSTRACT

Blood was taken from wild juvenile loggerhead sea turtles (Caretta caretta) that had been brought to the dock by cooperating fishermen during 1983-1985. Total protein, albumin, calcium, phosphorous, uric acid, creatinine, bilirubin, sodium, potassium, chloride, urea nitrogen, glucose, and corticosterone were measured in serum. Concentrations of these compounds changed in response to environmental changes in salinity, temperature and possibly prey availability. Profiles of juvenile loggerhead serum values were analysed using the multivariate methods of principal component analysis (PCA) and Q-Mode factor analysis. The results of PCA showed changes in sick animal's serum values attributable to impaired osmoregulation. The results of Q-mode factor analysis showed a strong relationship in animal groupings with temperature and salinity as well as stress. In most cases, stress effects were those associated with extreme pathological compromise or heat stress. Some cases showed no obvious correlation to any factor or variable measured and were attributed to underlying causes not detected in the survey methods. The resulting data provides background serum chemistry values for juvenile loggerhead sea turtles and illustrates the usefulness of the multivariate techniques of PCA and factor analysis in describing data relationships that are otherwise not obvious from general observation or univariate statistics.

ASSESSING ENVIRONMENTAL STRESS ON THE LOGGERHEAD SEA TURTLE (CARETTA  
CARETTA) IN VIRGINIA WATERS

## I. INTRODUCTION

Chesapeake Bay is a major foraging ground for immature Atlantic loggerhead (Caretta caretta) sea turtles (Lutcavage, 1981; Lutcavage and Musick, 1985). During each summer season sea turtle mortality increases drastically, beginning within 2-3 weeks after the arrival of turtles in the spring. Conservation and management personnel who are concerned with the long-term survival of sea turtles require an understanding of the role of environmental stress as a factor that influences the survival of individuals within regional populations of animals. The purpose of this work was to determine the value of serum components as a means to evaluate physiological and environmental stress on juvenile loggerhead turtles (C. caretta) in Chesapeake Bay.

Annual migration of sea turtles into Chesapeake Bay begins in late April and early May (Musick et al., 1984). Turtles reside in the Bay, foraging during the summer, and departing in early fall. Migration from the Bay corresponds with the first severe decline in water temperature, which generally occurs in October. Sea turtle mortality begins within 2-3 weeks after the arrival of turtles in the spring. Although mortality continues throughout the summer, it declines sharply at the end of June. This early summer mortality may comprise up to 78% of the mortalities observed during the entire year

(Lutcavage and Musick, 1985). Over 75% of the documented mortalities in Chesapeake Bay sea turtles since 1979 have occurred during May and June (Musick et al., 1984).

The cause of the mortality observed in Chesapeake Bay sea turtles could not be determined in over 66% of the cases examined for 1979-1984 (Musick et al., 1984; Bellmund et al., 1987). Known causes of mortality in other sea turtle populations stem from a variety of causes including: habitat destruction; predation of eggs, hatchlings, and adults; or interference from commercial fishing vessels (Hildebrand, 1981; Hillstad et al., 1981; Bacon, 1981; Ross, 1981). Possible causes for the mortality observed in Chesapeake Bay during May and June include natural population fluctuations, fishing gear entrapment, disease, and reduced condition due to over-wintering or migration.

Virginia Institute of Marine Science (VIMS) personnel determined the cause of death for 88 of 285 turtles examined during 1979-1983 (Musick et al., 1984, Lutcavage and Musick, 1985). Due to decomposition, most carcasses proved unsuitable for histological examination (Bellmund et al., 1987). Table 1 shows the relative percent for each of the 88 cases where the cause of death was determined. Extensive postmortem changes in beach stranded sea turtles precluded determination of the causes of death in these animals as is the case in most beach strandings of marine animals (Jacobson, personal communication; Stevens, personal communication).



TABLE 1.

Apparent Causes of Death in Virginia Sea  
Turtles Examined between 1979-1983  
(After Bellmund, et al., 1987)

Category	Number	Percent
Undetermined	197	69.0
Net Related	53	18.6
Shark Related	1	0.4
Prop Damage	21	7.4
Intentional (Human Induced)	9	3.2
Other Fishing Gear	4	1.4
Total	285	100

Many of the turtles that enter the Chesapeake Bay during spring are apparently in worse condition than those that depart in the Fall (Lutcavage and Musick, 1985). A hypothesis explaining this observation is that the first wave of migrants consists of animals that are healthy and animals that are diseased or stressed due to a variety of factors. Stressed or diseased animals die off early in the season and once these animals are removed from the population, mortality declines sharply. If this hypothesis is true then live animals arriving in the earliest portion of the season should constitute two groups with a range between two extremes. One group would survive the season and leave healthy in the fall and the other group would be pathologically compromised and die during the early part of the season.

Sources of stress during migration which may be responsible for mortality include; reduced feeding, energy expenditure, and temperature changes. Other stresses may result from previous capture and associated aspiration of seawater. Animals with these conditions may constitute the stressed or diseased portion of the population. This terminally stressed group would make up the portion of the population that die with no detectable causes in the early part of a summer season in Chesapeake Bay. Early season mortality may represent the death of sub-clinically ill animals.

Reptiles often show no visible signs of disease until actually becoming moribund (Frye, 1981; Lutz, 1972; Jacobson, personal communication). However, disease conditions may be reflected by

shifts in various blood and serum components prior to the animal becoming moribund. Blood has been examined in individuals in various wildlife populations in an attempt to evaluate the effects of environmental stresses (Grunow et al., 1984; Sturbaum and Bergman, 1981; Frye, 1981). These methods have been used to reflect the internal status of individual members of a population with no previous history on the individual. Lutz and Dunbar-Cooper (1987) used these methods to describe the physiological state of sea turtles from Florida waters. Sturbaum and Bergman (1981) used serum chemistry to describe changes induced by heat stress in the box turtle.

All animals that spend the summer in Chesapeake Bay experience the seasonal changes in physical regime associated with a temperate estuary. Resulting physiological changes are then due to factors which include the following: seasonal temperature fluctuations, salinity differences from area to area, and nutritional effects from variations in prey availability. In addition, other factors which may cause more drastic shifts in individual animal measurements include prolonged diving or increased activity during a tidal cycle (Lutcavage et al., 1987; Byles in prep.). These physical cycles affect physiological relationships and pathological conditions that are then reflected in serum chemistry of the general population of C. caretta in Chesapeake Bay.

The purpose of this study was to measure blood components, examine them in context of environmental changes present, and

evaluate them for use in describing the general pathological state of non-captive Virginia populations of juvenile loggerhead sea turtles. The parameters examined included enzymes, proteins, electrolytes, and nitrogenous waste products. Samples from animals were examined by location, broken into early season and late season, and grouped together for multi-variate analyses. Methods chosen to examine this data, beyond examination of mean trends, were principal component analysis (PCA) and Q-mode factor analysis. These procedures allow the examination of variability in blood components resulting from complex physiological and pathological interrelationships.

## II. METHODS AND MATERIALS

### Physical Examination and Location

Sea turtles used in this study were specimens that were incidentally captured alive in pound nets in the lower Chesapeake Bay. They were brought to the dock by pound net fishermen working in cooperation with personnel from VIMS. Pound nets are stationary fishing weirs, which provide free access to air in the head portion of the net and have large openings that allow free movement in and out of the nets for turtles (Lutcavage, 1981; Bellmund et al., 1987). Animals were kept in shade on the boat until their arrival at the dock and subsequent sampling (1-4hrs). Once at the dock, blood was removed using the methods described below, and the animals were examined and measured. After examination and sampling, animals were released from the dock. If the animal appeared stressed or it was not possible to sample blood at the dock, it was taken back to VIMS and allowed to recover in a tank with flowing sea water before blood was drawn. Locations were identified by zone (Figure 1). To facilitate other comparisons zones were combined to represent areas that may comprise homogeneous groups of animals. These areas and their respective zones are combined in Figure 2.

Figure 1. Zones for Virginia Study Site





Figure 2. Northern and Southern Areas For  
Blood comparisons





STUDY AREA

Weight and straight line length and width measurements were taken as described in Table 2. Dimensions of turtles were taken with one meter tree calipers to the nearest 0.1cm, and a tape measure to the nearest 0.5cm. Weight was taken with a spring balance to the nearest 0.5kg. Cloacal temperatures were taken, when possible, with a laboratory alcohol thermometer or a Schulteiss quick recording thermometer. Water temperatures and average salinities were taken from available Chesapeake Bay data.

#### Blood Sampling and Analysis

Blood was taken from live animals only. It was drawn aseptically from the cervical sinus using a non-heparinized 10cc syringe and a 20G:1 1/2 needle (Owens and Ruiz, 1980; Bentley and Dunbar-Cooper, 1980). Blood was transferred to a siliconized 10cc vacutainer tube and allowed to clot at 4°C for 45 minutes. Blood for serum chemistry was spun at 6000g for 25 mins. Serum was removed with a pipet and transferred into a separate test tube for storage. Serum samples were frozen at -20°C to -40°C until analysis.

Serum was analyzed by several techniques. Testosterone and corticosterone were determined using radioimmune assay (RIA) at Texas A&M University (Morris, 1982). Hormones were measured using RIA techniques according to Owens (1978), Morris (1982), and Morris and Owens (1982).

Table 2.  
Sea Turtle Morphometric Measurements

Measurement	Description
Straightline carapace length (CLS)	Length of top shell at longest points, taken with calipers.
Straightline carapace width (CWS)	Width taken at widest point with calipers.
Curved carapace length (CLC)	Longest length over curve using a tape measure.
Curved carapace width (CWC)	Width at widest point, over curve using a tape measure.
Plastron length (PL)	Length of plastron at longest point.
Plastron width (PW)	Width at widest point of plastron, not including bridge plates, taken with calipers.
Plastron with bridge (PWB)	Width at widest point, includes bridge plates, taken with calipers.
Weight	Taken with spring balance to nearest 0.5 kg.

All sera were analyzed using a Hitachi 705 blood analyzer system set to measure reactions within the linear range of sample concentrations (Appendix I). Serum enzyme levels were too low or too variable in concentration to measure due to sample handling effects rather than machine range.

Sera was taken from one individual animal held at two constant temperatures. Blood was sampled before and after feeding, over time, to monitor the change in glucose and urea nitrogen concentrations (Appendix II). Blood was taken once before feeding and sequentially in twenty minute intervals for the first two hours and then approximately every two hours until concentrations decreased to original pre-feeding levels. The first temperature used was 20°C and the animal had been held at that temperature for several months. The animals body temperature was then raised to 25°C over a 4 day period. The animal was not allowed time to physiologically acclimate and the same procedures were then carried out at this temperature. Prior to the first feeding, food was with-held from the animal for seven days. The animal was given 6.6kgs of thawed striped bass (Morone saxitilis) for both feedings. The turtle, was a small male weighting 32 kg with shell length, CLS of 59.4 cm and width, CWS of 48.5 cm.

Blood was taken from live animals only. Serum concentrations of the following constituents were measured; total protein in g/dl (TPRO), albumin in g/dl (ALB), globulins in g/dl (calculated as difference from TPRO - ALB), serum glutamate oxaloacetate

transaminase (SGOT), lactate dehydrogenase (LDH), alkaline phosphatase (ALK), calcium in mg/dl (Ca), inorganic phosphorus in mg/dl (Pi), sodium in MEQ/l (Na), chloride MEQ/l (Cl), potassium in MEQ/l (K), creatinine in mg/dl (Crea), uric acid in mg/dl (UA), blood urea nitrogen (BUN) in mg/dl, total bilirubin, glucose in mg/dl (glu), testosterone in ng/ml, and corticosterone in pg/ml (cort). The condition factor  $Q$  (where  $Q = \text{weight}/(\text{length})^3$ ), proportional to density of the animal, was calculated and used when appropriate. Turtles that were classified as being apparently healthy were used for all mean calculations unless noted. Means were calculated using a single sample per individual, no double samples were included unless noted.

Initially, results were examined for mean trends with means given as  $\pm$  SD. For comparison and statistical analyses, animals were classified by sex, month, size, location and state of health. Sex was determined by testosterone titer in serum. Animals were grouped into two size categories, those  $< 70.0$  cm and those  $> 75.0$  cm, using 75.0 cm CLS as the lower size limit for nesting females (Martin, personal communication; Richardson *et al.*, 1978). Turtles 70-75 cm were used as dividing groups and were not included in this Table only. Samples for individual animals were divided by location as previously described. Turtles were subjectively defined as apparently (or clinically) healthy if they were not emaciated, injured or obviously ill, otherwise they were defined as sick.

### Multivariate Analysis

Multivariate analysis was evaluated to determine effectiveness in relating internal homeostasis to health using blood values from field sampled animals. In this work, the multivariate techniques of principle component analysis (PCA) and Q-mode factor analysis were used to examine relationships in serum components and individual animals. Principle component analysis was run using the 'Factor' routine without iteration on a correlation table in Statistical Package for the Social Sciences (SPSS) Release 7-9 (Nie et al., 1975; Hull and Nie, 1981). Homogeneity of variance was tested using Cochran's test and Bartlett-Box F in the 'Manova' routine of SPSS release 7-9 (Nie et al., 1975; Hull and Nie, 1981). All multivariate analyses were run on object sets with no missing variables. Q-mode analyses were run using CABFAC, QMODEL, and EXQMODEL, programs developed by Klován and Miesch (1975). Their modification and adaptation to the PRIME are discussed by Berquist (1986).

#### Factor Analysis

Factor analysis is a multivariate technique essentially resulting in a reduction of variable space. Its analytical intent is to find  $m$  linear combinations of original  $n$  variables to describe real entities without significant information loss (Klován, 1975). An in-depth discussion of factor analytic techniques can be found in Davis (1986) and Joreskog, et al. (1976). The following discussion is a general description of these methods as they were applied to this data set and relies heavily on these texts.

Factor analytic methods can be divided into two general techniques. These techniques are; R-mode, which includes Principal Component Analysis (PCA), and Q-mode. R-mode techniques place variables in groups and measure the interrelationships between these groups of variables. Q-mode techniques measure interrelationships between objects (individual turtles sampled). Differences between PCA and 'true' R-mode analyses are due to factor determination methods and lack of a statistical model for PCA (Joreskog et al., 1976). In component analysis factors are defined to maximize the percent variance explained. In 'true' R-mode factor analysis factors are chosen to account for the maximum amount of intercorrelation of the variables (Joreskog et al., 1976). Functionally R-mode analyses describe interrelationships within a data matrix of  $m$  variables and reflect the correlation of each variable with  $p$  mutually uncorrelated underlying factors, assuming  $p < m$ . (Davis, 1986).

#### Principal Components

Principal components are the eigenvectors of a variance-covariance or correlation matrix, depending upon the original calculations. In principal component analysis the maximum amount of variance will be resolved by the first principal axis. If a matrix is represented as a series of vectors in multidimensional space and each row of the matrix give the coordinates of the vector represented by that row, then, as in the  $2 \times 2$  example, the vectors define arbitrary axes of a hyperdimensional figure. The eigenvectors now define the orientation of the principal axes of the hyperdimensional



figure and the eigenvalues represent the magnitude of the variance explained by each principle axis. In this way the linear combination of variables that explains the maximum amount of variation in the data can be found. PCA in the SPSS routine bases calculations on the correlation matrix and so resulting groups are unitless (Joreskog et al., 1976).

PCA was then run on groups arranged by time from the apparently healthy animals. Animals were divided into early season (May and June) and late season (July, August, and September). Relationships in this breakdown by time were indeterminant so the data was then analyzed using Q-mode factor analysis.

#### Q-Mode Analysis

Q-mode analysis was run on all single Caretta blood value sets that had no missing values. The Cosine theta similarity coefficient index was used to assess relatedness. The following serum parameters were used in this analysis;

Total Protein	Sodium
Albumin	Potassium
Calcium	Chloride
Phosphorus	Blood Urea Nitrogen
Uric Acid	Glucose
Creatinine	



### III. RESULTS

#### Preliminary evaluation:

Means values of blood serum parameters were calculated by month (Table 3), by location (Table 4), by sex (Table 5), by size (Table 6), and by condition (Table 7). Means are not reported for serum enzymes since they were found to be extremely variable and were affected by differences in handling of samples. Means for uric acid and creatinine are included but the interpretation of these values is uncertain since circulating concentrations of these compounds are at or below the detection levels of the analytical methods used. Means for proteins (Figure 3), serum ions (Figure 4), and organics (Figure 5) are plotted by month for all years combined.

The northern-most site (Potomac River Mouth) represented a discrete location with enough animals to be examined separately (N=43 apparently healthy animals). Mean values for turtles captured from this site were then compared to values derived from the combined data collected from the southern sites. The distance (42.6 km) between the Potomac River site and the most northerly of the combined sites (Cherry Point) is great enough difference to allow this grouping of samples. These sites were divided into zones by Lutcavage (1981) and zones were used to maintain consistency with previous samples from Virginia waters. Zones as shown in Figure 1 and are grouped into

Table 3.  
Healthy Caretta caretta Serum Values for Summer Seasons 1983-1985

Month	Parameter	TPRO	Alb	Ca	P:	UA	Creat.	Na	K	Cl	BUN	GLUC	Na/K	K/BUN	Na/BUN
May	mean	3.4	1.1	7.1	4.6	1.4	0.2	161	3.6	112	48	103	46.3	8.6	38.7
	SD	0.7	0.3	3	1.3	0.6	0.1	3.6	0.6	5.6	18.9	28.3	7.9	3.8	14.5
	N	12	12	12	12	11	12	12	12	12	12	12	12	12	12
June	range	2.7-4.6	0.7-1.9	2.2-10.4	3.1-7.1	0.7-2.2	0.1-0.3	155-168	2.7-4.3	103-122	26-78	72-161	39-60	3.7-15.5	20.1-63.0
	mean	3.3	1.2	4.6	5.1	1.1	0.2	159	3.9	113	82	159	41.5	5.0	20.7
	SD	0.7	0.4	2.7	1.4	0.4	0.1	6.4	0.6	7.5	19.3	64.9	6.0	1.4	6.4
July	N	31	31	31	31	31	31	31	31	31	31	31	31	31	31
	range	2.0-4.5	0.6-2.0	1.4-9.7	2.3-7.7	0.2-2.2	0.1-0.4	140-167	2.9-4.9	90-123	40-108	22-292	33-55	3.1-8.8	14.6-39.7
August	mean	3.5	1.3	5.5	6.0	0.9	0.3	158	4.7	110	85	170	34.5	6.2	21.3
	SD	0.7	0.4	2.2	1.4	0.4	0.1	6.3	0.8	6.7	23.2	84.6	5.5	3.2	12.3
	N	16	16	10	10	16	16	16	16	16	16	16	16	16	16
September	range	2.1-5.0	0.8-2.5	2.3-8.9	2.5-8.5	0.2-1.5	0.2-0.5	143-168	3.4-6.0	97-122	24-124	81-458	26-45	4.2-17.5	12.6-68.8
	mean	3.4	1.3	5.3	5.2	1.2	0.3	153	4.8	110	106	184	32.3	4.7	15.2
	SD	0.6	0.4	2.4	1.2	0.3	0.1	10.7	0.7	7.9	18.6	46.3	4.9	1.2	4.6
October	N	6	6	6	6	6	6	6	6	6	6	6	6	6	6
	range	2.7-4.4	0.8-2.0	3.2-9.6	3.7-7.0	0.7-1.6	0.2-0.4	136-166	3.9-6.1	95-119	68-115	113-243	25.3-38.2	3.4-6.8	12.3-24.4
November	mean	3.8	1.5	5.5	6.8	0.7	0.3	153	3.5	110	67	99	44.5	5.9	25.8
	SD	0.5	0.2	2.3	1.6	0.1	0.1	8.4	0.6	6.5	25.1	39.7	6.9	2.6	10.9
	N	5	5	5	5	5	5	5	5	5	5	5	5	5	5
December	range	3.0-4.4	1.3-1.8	2.1-7.7	5.2-8.8	0.5-0.8	0.2-0.3	145-164	2.6-4.0	101-118	36-105	70-167	38-56	3.7-10.3	15.6-44.2

Table 4.  
Healthy Caretta caretta Serum Values by Zone 1983-1985

Zone	Parameter	TPRO	Alb	Ca	P:	UA	Creat.	Na	K	Cl	BUN	GLUC	Na/K	KBUN	Na/BUN
Zone 3	mean	3.1	1.0	2.9	3.9	1.0	0.2	160	3.6	117	73	113	45.3	5.6	25
	SD	0.6	0.4	1.2	1.5	0.5	0.1	4.4	0.7	4.3	30.0	54	7.0	1.9	10
	N	8	8	7	8	8	8	8	8	8	8	8	8	8	8
	range	2.5-4.2	0.6-1.6	2.0-5.5	2.3-5.5	0.6-2.2	.01-0.4	155-167	3.1-4.9	109-121	40-108	22-197	34-51	3.1-8.9	14.6-38.7
Zone 7	mean	3.5	1.3	4.5	5.4	0.7	0.26	158	3.8	112	74	134	42	7.2	28.6
	SD	0.4	0.3	2.3	1.5	0.3	0.1	7.7	0.4	7.0	34	64	3.1	5.9	21.3
	N	5	5	5	5	5	5	5	5	5	5	5	5	5	5
	range	2.9-5.0	1.0-1.7	1.9-7.9	3.8-7.3	0.2-1.0	0.2-0.3	145-164	3.3-4.2	101-117	24-107	86-246	38-45	3.5-17.5	15-65.8
Zone 9	mean	3.3	1.1	5.7	5.2	0.9	0.2	156	3.5	114	62	114	46	6.0	27.6
	SD	0.5	3.3	2.8	1.8	0.6	0.1	7.5	0.6	5.6	19.4	41	7.6	2.0	9.6
	N	9	9	9	9	9	9	9	9	9	9	9	9	9	9
	range	2.7-4.1	0.7-1.6	1.4-8.7	3.1-8.0	0.2-2.2	0.1-0.4	146-168	2.6-4.3	107-122	36-97	73-197	28-59	2.7-10.3	16-44.2
Zone 12	mean	3.6	1.3	5.7	5.7	1.1	0.3	158.2	4.3	110.2	83	176	38	5.5	20.5
	SD	0.6	0.4	2.8	1.3	0.4	0.1	6.9	0.8	6.6	21	66	7.4	1.4	6.8
	N	43	43	43	43	43	43	43	43	43	43	43	43	43	43
	range	2.1-5.0	0.7-2.5	2.1-10.5	3.1-8.5	0.5-2.0	0.1-0.5	136-168	2.7-6.1	90.0-123	37-124	77-458	25-60	3.4-9.7	12-42

Table 5.  
Serum Values for Healthy Caretta caretta by Sex 1983-1985

	Parameter	TPRO	Alb	Ca	P:	UA	Creat.	Na	K	Cl	BUN	GLUC	Na/K	K/BUN	Na/BUN
Females	mean	3.2	1.0	3.2	4.7	1.0	0.3	157	4.2	113	82	146	39	5.5	21.7
	SD	0.5	0.3	1.1	1.4	0.4	0.1	7.3	0.8	7.1	26	55	8.0	1.6	8.4
	N	21	21	21	21	21	21	21	21	21	21	21	21	21	21
	range	2.5-4.5	0.6-2.0	1.9-5.5	2.3-7.0	0.2-2.2	0.1-0.4	136-167	2.7-5.8	95-123	41-114	22-246	27-58	3.1-8.8	12.3-38.3
Males	mean	4.0	1.4	3.9	6.2	1.0	0.3	159	4.3	113	76	149	38	6.0	23.5
	SD	0.5	0.3	1.3	1.7	0.3	0.1	6.9	0.9	4.9	27	49	8.4	1.3	8.9
	N	7	7	7	7	7	7	7	7	7	7	7	7	7	7
	range	3.5-5.0	0.9-1.6	2.7-5.4	3.2-8.2	0.6-1.4	0.1-0.4	149-168	3.1-5.2	108-122	40-114	70-207	365-50.0	4.1-7.7	13.4-38.7

Table 6.  
Serum Values for Healthy Caretta caretta by Size 1983-1985

Size	Parameter	TPRO	Alb	Ca	P:	UA	Creat.	Na	K	Cl	BUN	GLUC	Na/K	K/BUN	Na/BUN
CLS <70.0 cm	mean	3.3	1.2	4.8	5.3	1.0	0.2	157	4.0	112	80	143	40.5	5.6	22.7
	SD	0.6	0.3	2.5	1.5	0.4	0.1	7.3	0.8	7.5	26	52	8.1	2.7	11.3
	N	49	19	45	49	19	48	49	49	49	49	49	49	49	49
	range	2.0-4.6	0.6-2.0	1.4-10.4	2.3-8.2	0.2-2.2	0.1-0.4	136-168	2.6-6.0	90-123	24-124	22-246	26.2-60.4	3.1-17.5	12.2-65.8
CLS >75.0 cm	mean	3.9	1.4	7.3	5.6	1.3	0.2	161	4.2	113	68	167	40.3	7.2	30.3
	SD	0.5	0.4	2.6	1.4	0.6	0.1	4.4	0.9	5.7	28	100	9.7	3.1	18.7
	N	18	18	16	18	18	18	18	18	18	18	18	18	18	18
	range	2.9-5.0	0.7-2.5	2.1-10.3	3.2-8.8	0.5-2.7	0.1-0.5	150-168	2.3-6.1	97-122	20-115	57-458	25.2-70.8	3.8-13.8	13.4-81.5

Table 7.  
Serum Values for Caretta caretta 1983-1985

	Parameter	TPRO	Alb	Ca	P:	UA	Creat.	Na	K	Cl	BUN	GLUC	Na/K	K/BUN	Na/BUN
Apparently Healthy	mean	3.5	1.2	5.5	5.4	1.2	0.2	158	4.0	112	76	147	40	6.0	24.6
	SD	0.7	0.4	2.7	1.5	1.2	0.1	6.8	0.8	6.9	27	68	8.3	2.8	13.5
	N	75	75	69	75	75	73	74	74	74	75	75	74	74	74
	range	2.0-5.0	0.6-1.9	1.4-10.4	2.3-8.8	2.0-11.0	0.1-0.5	136-168	2.3-6.1	90-123	9-124	17-458	25-71	3.1-17.5	12.3-81.5
Sick or Injured	mean	3.9	1.8	6.5	6.3	1.2	0.3	150	3.5	108	54	138	47.4	8.4	37.9
	SD	1.2	0.6	2.0	2.0	1.0	0.1	16.2	1.3	14.9	24	67	14.2	4.8	22.7
	N	14	14	14	14	14	14	12	12	12	14	13	12	12	12
	range	1.6-6.1	0.3-2.7	2.4-8.9	3.3-11.5	0.3-4.3	0.1-0.4	105-168	2.1-6.7	64-124	19-87	67-239	15.6-70.5	3.7-15.5	17.5-77.9

Figure 3.  
Mean Serum Protein Values for All Healthy Animals




Figure 4.  
Mean Serum Electrolytes for All Healthy Turtles







Figure 5.  
Serum Organics for all Healthy Turtles



areas in Figure 2. Mean concentrations were plotted for the Potomac River site (Area 4) and compared with southern sites (Area 1,2,3: Zones 3, 5, 6, 7, and 9) for proteins (Figure 6), serum ions (Figure 7), and organic components (Figure 8).

#### Total Protein

Mean values for total protein ranged from 3.3 - 3.8 g/dl over the season. Mean total protein values for males were  $4.0 \pm 0.5$  g/dl while those of females were  $3.2 \pm 0.5$  g/dl. Mean total protein values were  $3.3 \pm 0.6$  g/dl for animals less than 70.0cm CLS and  $3.9 \pm 0.5$  g/dl for turtles greater than 75.0cm CLS. Apparently healthy turtles had a mean total protein of  $3.5 \pm 0.7$  g/dl, while sick or injured animals had a mean value of  $3.9 \pm 1.2$  g/dl.

#### Albumin

Although, mean albumin values increased steadily from  $1.1 \pm 0.3$  g/dl to  $1.5 \pm 0.2$  g/dl from May to September there was no pattern to mean values from different locations. Mean albumin values were  $1.0 \pm 0.3$  g/dl for females and  $1.4 \pm 0.3$  g/dl for males. The mean albumin concentration was  $1.2 \pm 0.3$  g/dl for animals less than 70.0cm CLS and  $1.4 \pm 0.4$  g/dl for animals greater than 75.0cm CLS. The mean value for apparently healthy animals was  $1.2 \pm 0.4$  g/dl while the mean value for this constituent was  $1.8 \pm 0.6$  g/dl for sick or injured animals.

#### Calcium

Calcium values were highest in May ( $7.1 \pm 3.0$  mg/dl) and October ( $7.3 \pm 2.1$  mg/dl), while June through September ranged from 4.6 ( $\pm$

Figure 6.  
Seasonal Differences in Proteins of *Caretta*  
from Different Regions 1983-1985




Figure 7.  
Seasonal Differences in Electrolytes of *Caretta* from  
Different Regions 1983-1985





Figure 8.  
Seasonal Differences in Organic Components  
from *Caretta* of Different Regions 1983-1985





2.7) to 5.5 ( $\pm 2.3$ ) mg/dl, respectively. Calcium levels were  $3.2 \pm 1.1$  mg/dl in females and  $3.9 \pm 1.3$  mg/dl in males. Serum calcium was  $4.8 \pm 2.5$  mg/dl in animals less 70.0 cm and  $7.3 \pm 2.6$  mg/dl in animals greater than 75cm.

#### Phosphorus


When plotted over the season, there was a trend towards increasing serum inorganic phosphorus concentration during the summer. When plotted together phosphorus values followed calcium values closely in healthy animals (Figure 9). Phosphorus values were  $6.2 \pm 1.7$  mg/dl in male turtles and  $4.7 \pm 1.4$  mg/dl in females. Mean phosphorous was  $5.4 \pm 1.5$  mg/dl in apparently healthy animals and  $6.3 \pm 2.0$  mg/dl in sick and injured turtles.

#### Sodium

This was the most variable electrolyte from area to area. Electrolytic differences were most apparent in this parameter. Differences were most visible when this parameter was examined separately for the northern-most site and this data compared with the most extreme southern site. Although mean values were within 2 meq/l for all areas, standard deviations were less ( $sd=4.4$ ,  $n=8$ ) in the high salinity Bay mouth site (Zone 3) than in the lower salinity northern River site ( $sd=6.9$ ,  $n=43$ ). Animals over the season only ranged from 151 to 164 in the northern river site (Figure 7), while animals in the southern (Zone 3) site ranged from 149 to 163 (Figure 7). Mean sodium values were more variable in smaller animals (less than 70.0 cm),  $sd=7.3$ ,  $n=49$ , while larger animals (greater than 75.0cms),  $sd=4.4$ ,  $n=18$  showed less variation in this parameter. Mean

Figure 9. Seasonal Calcium and  
Phosphorus Concentrations

# Mean Seasonal Calcium and Phosphorus Concentrations





sodium values showed an inverse relationship with albumin(Figure 10). Mean values for apparently healthy animals was  $158 \pm 6.8$  meq/l and  $150 \pm 16.2$  meq/l in sick or injured turtles.

#### Chloride

Mean chloride concentration decreased from a high value of  $117 \pm 4.3$  meq/l in the Bay mouth site, to a low of  $110 \pm 6.6$  meq/l in the Potomac River site. Chloride values changed with sodium over the season (Figure 11). Chloride and sodium were tightly regulated in C. caretta. Ranges for this parameter were greatest in animals from the low salinity Potomac River site, Cl ranged from 90-123 meq/l in this area (Zone 12). Sick turtles showed the greatest variation in this component. Apparently healthy turtles had chloride values ranging from 90-123 meq/l with a mean of  $108 \pm 6.9$  meq/l, while sick turtles had a range of 64-124 meq/l with a mean of  $108 \pm 14.9$  meq/l.

#### Potassium

Potassium values for animals from the Potomac River (Zone 12) had a wide range (2.7-6.1meq/l) while the range for the Bay mouth animals was 3.1-4.9 meq/l. When values from zone 12 were plotted by month and compared to those values from all other areas it showed a steady increase through most of the summer, while other areas plotted separately showed a more random pattern (Figure 7). Mean values for this electrolyte increased during the summer (Figure 7). Mean values for this electrolyte were similar in both male ( $4.3 \pm 0.9$  meq/l) and female turtles ( $4.2 \pm 0.8$  meq/l) (Table 5). Apparently healthy animals had a mean value of  $4.0 \pm 0.8$  meq/l, while sick and injured turtles had a mean value of  $3.5 \pm 1.3$  meq/l. This parameter

Figure 10. Seasonal Sodium and Albumin  
Concentrations

# MEAN NA AND ALB CONCENTRATION BY MONTH





Figure 11. Mean Sodium and Chloride  
Concentration by Month

# MEAN NA AND CL CONCENTRATION BY MONTH



exhibited a direct relationship with urea nitrogen in healthy animals (Figure 12).

#### Blood Urea Nitrogen

Blood urea nitrogen increased from a low mean value of 48 mg/dl in May to a peak mean value of 106 mg/dl in August. Apparently healthy animals had mean values of  $76 \pm 27$  mg/dl, while sick or injured animals had a mean value of  $54 \pm 24$  mg/dl. This parameter was variable in all cases and had consistently wide ranges.

#### Glucose

Potomac River turtles showed the widest range and had higher mean levels of glucose than turtles from any of the other sites. Glucose increased over the summer through August, and declined in September and October in both sick and healthy animals. The mean and standard deviations were greater in large turtles, but were not affected by sex. Levels of urea nitrogen and potassium mirrored this temporal and spatial increase in mean glucose values (Figure 13). Glucose was variable and showed large deviations in healthy animals, large animals, animals collected during July, and animals collected from Zone 12.

#### Glucose and Urea Nitrogen Analysis

Results of preliminary examination of the changes in glucose-urea nitrogen as a function of temperature are shown in Figure 14. Urea nitrogen rose at about the same time at both temperatures. Glucose remained elevated longer at 26° C than at 20° C. Since the animal was not allowed time to acclimate physiologically, this can be

# MEAN NA AND CL CONCENTRATION BY MONTH




Figure 12. Mean Seasonal Potassium and  
Urea Nitrogen Concentrations



# Mean Seasonal Potassium and Urea Nitrogen Concentrations




Figure 13. Seasonal Values for Potassium

MEAN SERUM POTASSIUM  
FOR 1983-1985






Figure 14. Glucose and Urea Nitrogen  
Response to Temperature

# Glucose and Urea Nitrogen Response to Temperature

GLUCOSE AND BUN RESPONSE  
WITH TIME AT 20°C



GLUCOSE AND BUN RESPONSE  
WITH TIME AT 26°C



assumed to be a physical response to temperature only. Glucose may have had a more uniform response at the higher temperature but more analyses on more animals must be done to make any conclusive statements about these relationships.

#### Corticosterone

Mean corticosterone values were plotted seasonally for 1984-1985 (Figure 15). Values were not available for animals from 1983. Multiple sample runs for this analyses were not consistent, probably due to the presence of an interfering substance. However, this compound was not found in all samples, was not indicated in the standards, or the recovery for the analysis and was never identified. Values for this hormone are therefore used as a general indicator only.

#### Multivariate Analysis


##### Principal Components Analysis

Principal component analysis (PCA) of serum components yielded general relationships which differed between healthy and sick turtles. The three major factor groups determined for healthy animals were:

<u>Group 1</u>	<u>Group 2</u>	<u>Group 3</u>
Albumin	Urea Nitrogen	Sodium
Total Protein	Potassium	Chloride
Calcium	Glucose	
Phosphorus		

Figure 15. Mean Serum Corticosterone for  
1984-1985

MEAN SERUM CORTICOSTERONE  
FOR 1984-1985





When serum values from the sick and injured group of animals were then analyzed, variable groupings changed. Mean values for those animals classified as sick or injured are found in Table 7. Variable groups for sick animals were:

<u>Group 1</u>	<u>Group 2</u>	<u>Group 3</u>
Albumin	Sodium	Urea Nitrogen
Total Protein	Chloride	Glucose
Calcium	Potassium	
Phosphorous		

The first group was composed of the same factors in both sick and healthy animals. Factors in the second group changed so that this grouping for sick animals included potassium, sodium, and chloride. The third group for sick animals was then composed of urea nitrogen and glucose. Potassium moved from the urea nitrogen/glucose group in healthy animals to the electrolyte group in apparently sick animals. The entire electrolyte group moved from third most important factor in healthy animals to second in sick and injured animals.

#### Q-Mode Analysis

General statistics for all turtles included in this analysis are found in Table 8. The analysis was run on 11 variables and 74 samples for 99% error explanation. Using the varimax solution, two factors explained 99.01% variance (dissimilarity) of the data with the relative input from each factor being 51.638% and 47.371%, respectively. The varimax factor loadings are listed in Table 9. The data best fit a two factor model. Best fit to a factor model was

Table 8.  
Mean Sera Values for Samples Included  
in Multi-Variate Analysis

Variable	No.	Mean	Standard Deviation	Minimum Value	Maximum Value
TPRO	1	3.5	0.78	1.6	6.1
ALB	2	1.3	0.47	0.3	2.7
CA	3	5.6	2.66	1.4	10.4
PHOS	4	5.5	1.53	2.3	11.5
UA	5	1.1	0.6	0.2	4.3
CRE	6	0.2	0.1	0.1	0.5
NA	7	157	9.4	105	168
K	8	3.9	0.89	2.1	6.7
CL	9	111	8.6	64	123
BUN	10	72	27	19	124
GLUC	11	150	68.4	22	458

Table 9.  
 Program CABFAC Results of the VARIMAX  
 Factor Score Matrix

Variables	No.	Factor 1	Factor 2
Total Protein	1	0.016	0.004
Albumin	2	0.004	0.003
Calcium	3	0.027	0.004
Phosphorus	4	0.021	0.010
Uric Acid	5	0.005	0.001
Creatinine	6	0.000	0.001
Sodium	7	0.776	0.101
Potassium	8	0.013	0.009
Chloride	9	0.566	0.054
Blood Urea Nitrogen	10	0.230	0.173
Glucose	11	-0.152	0.978

based on evenness of the variance explained. The addition of factors did not evenly divide variance, further factors only accounted for 3-4% of variance. Factor one loaded heavily on sodium, chloride, urea nitrogen, and negatively on glucose.

Solutions showed factors to be made up of two groups. Endmembers chosen using Imbries solution were MT-21-84 for Factor 1 and MT-88-83 for Factor 2. Using Imbries solutions all turtles were plotted and are shown in Figure 16. Animals grouped at either extreme (endmember turtle groups) are marked in Figure 17. Collective data for endmember turtles from Factor Group 1 are listed in Table 10. Collective data for endmember turtles from Factor Group 2 are found in Table 11. In addition to endmember animals, other animals grouped together with factor scores of .7900-.8100. Animals falling into this intermediate group collective data is provided in Table 12. Although this group shows no obvious direct trends, it contains two animals that were sub-clinically ill, (MT-12-84) and (MT-124-84). This group contains several stressed animals (MT-36-83, MT-32-84, and MT-62-83).

The compositions of the Factor groups are controlled by the variable loadings. Factor one loaded heavily on sodium, chloride, and urea nitrogen. Factor two loaded on glucose, urea nitrogen, and sodium, however input from glucose was much higher than that of the other two variables. The composition of Factor group one was controlled by sodium and chloride with some input from urea nitrogen while the composition of factor group two was controlled by glucose

Figure 16. Turtle Grouped by Serum Variables:  
Varimax Factor Loadings for a Two  
Factor Solution

Turtles Grouped by Serum Variables:  
Varimax Factor Loadings for a Two Factor Solution




Figure 17. Turtles Grouped by Serum  
Variables: Varimax Factor Loadings  
for a Two Factor Solution

Turtles Grouped by Serum Variables:  
Varimax Factor Loadings for a Two Factor Solution






Table 10 . Endmember Groups and Serum Profiles for Factor Group 1

ID no.	Marine Turtle No.	Sam-ple No.	Factor Score	Date	Tpro (g/dl)	alb (g/dl)	Glob (g/dl)	Ca (mg/dl)	Phos (mg/dl)	Uric acid (mg/dl)	Cre (mg/dl)	Na (meq/l)	K (meq/l)	Na/K	Cl (meq/l)	BUN (mg/dl)	Glu (mg/dl)	Wt (kg)	CLS (cm)	CWS (cm)	Zone	Comments
19	MT-21-84	5757	.9655	1-VI-84	2.6	0.6	2.0	2.3	3.4	2.2	0.1	159	3.2	49.7	121	57	22	22	53.7	45.6	3	From Lynnhaven, high salinity apparently healthy. early season cool water animal
55	MT-140-84	5809	.9014	12-XII-84	4.2	1.2	3.0	8.0	4.6	2.7	0.1	163	2.3	70.9	119	20	57	57.5	75.7	61.4		Cold stunned, from Eastern Shore of VA. From 3°C water, 8°C after 4hrs, 14°C when sampled 12hrs after arrival, apparently healthy
57	MT-04-85	5853	.8665	9-V-85	4.4	1.4	3.0	10.3	5.2	1.4	0.2	164	3.6	45.5	116	26	76	120.0	89.9	72.9	n/a	Taken offshore of VA BCH during trawl trip, apparently healthy, cloacal temp. 17°C.
50	MT-129-84	5800	.8743	22-IX-84	4.2	1.5	2.7	5.4	8.2	0.8	0.2	149	3.9	38.2	109	65	70	47.0	67.6	55.4	n/a	Taken in Norfolk Canyon during longline cruise. Cloacal temp. 20°C.
66	MT-48-85	5866	.8648	9-VI-85	4.4	1.5	2.9	7.7	5.6	0.7	0.2	162	2.9	55.8	109	80	77	45.0	66.4	54.9	12	Turtle from Potomac River, held at VIMS, this sample from one month after arrival. Apparently healthy
56	MT-03-85	5852	.8522	9-V-85	3.3	1.0	2.3	8.5	4.3	2.1	0.1	164	4.2	39	116	31	84	101.0	89.3	70.7	n/a	Taken on Offshore trawl trip, off VA BCH. Cloacal temp. 18°C. Apparently healthy.

Table 11 . Endmember Groups and Serum Profiles for Factor Group 2

ID no.	Marine Turtle No.	Sam- ple No.	Factor Score	Date	Tpro (g/dl)	alb (g/dl)	Glob (g/dl)	Ca (mg/dl)	Phos (mg/dl)	Uric acid (mg/dl)	Cre (mg/dl)	Na (meq/l)	K (meq/l)	Na/K	Cl (meq/l)	BUN (mg/dl)	Glu (mg/dl)	Wt (kg)	CLS (cm)	CWS (cm)	Zone	Comments
8	MT-88-83	5568	.9635	2-VIII-83	4.2	2.5	1.7	8.9	8.5	1.0	0.5	162	5.9	27	97	85	458	68.0	72.2	62.3	12	Stressed, fought prior to sample difficult to get blood, hot
12	MT-113-83	5578	.9271	7-X-83	4.0	1.9	2.1	8.3	6.0	4.4	0.4	106	7.0	15	59	57	244	36.0	65.1	55.1	15	Sick turtle from the Eastern shore of VA (Nassawadox Creek) died shortly after sampling, severely emaciated
69	MT-54-85	5872	.8868	19-VI-85	3.7	1.5	2.2	9.7	7.0	1.3	0.3	164	4.5	36	116	81	292	86.5	83.7	66.9	12	Recapture from 1984, apparently healthy when sampled
34	MT-83-84	5773	.8623	22-VI-84	4.5	2.0	2.5	4.9	4.9	1.0	0.3	154	4.3	35.8	109	53	246	31.0	55.6	50.0	12	Apparently healthy, from Potomac River, sampled at dock and released
13	MT-119-83	5577	.8525	29-X-83	3.5	1.6	1.9	6.6	4.9	1.1	0.2	150	2.4	62.5	112	29	239	16.0	49.1	43.7	7	Turtle with respiratory problems that died in captivity after apparent recovery. Histo. and necropsy showed extensive lung damage with systemic infection. Taken from Poquoson VA
14	MT-120-83	5572	.8438	2-XI-83	3.5	2.0	1.5	8.3	5.2	0.9	0.2	148	2.1	70.4	113	19	233	31.0	62.7	52.6	3	Taken from Portsmouth VA, emaciated severely lethargic- leeches on neck and head. Died 5-VIII-84.
65	MT-43-85	5868	.8534	6-VI-85	4.1	1.5	2.6	9.7	6.4	0.9	0.2	161	4.4	36.6	112	80	247	49.0	70.0	59.3	12	Apparently healthy heat stressed cloacal temp. 30°C taken from Potomac River
45	MT-120-84	5793	.8481	13-VIII-84	3.9	1.5	2.4	6.5	5.5	1.6	0.4	154	6.1	25.2	111	115	243	87.0	82.1	67.0	12	Potomac River turtle, sampled at the dock, apparently healthy, cloacal temp. 33.5°C

Table 12 . Intermediate Turtles Described by Factor Analysis

ID no.	Marine Turtle No.	Sam-ple No.	Date	Factor Score	Tpro (g/dl)	alb (g/dl)	glob (g/dl)	Ca (mg/dl)	Phos (mg/dl)	Uric acid (mg/dl)	Cre (mg/dl)	Na (meq/l)	K (meq/l)	Na/K	Cl (meq/l)	BUN (mg/dl)	Glu (mg/dl)	Wt (kg)	CLS (cm)	CWS (cm)	Zone	Comments
2	MT-36-83	5591	6-VI-83	.8093	2.4	0.8	1.6	6.2	3.0	1.7	0.2	144	3.4	42.4	91.0	96	85	28	58.0	50.0	5	Taken 6-IV-83 in pound net off Grandview apparently healthy, heat stressed.
5	MT-78-83	5597	12-VII-83	.7906	3.6	1.7	1.9	7.9	7.3	1.0	0.2	158	4.2	37.6	109	24	105	37	64.5	53.2	7	Taken from pound nets, York River mouth, held at lab, apparently healthy. Taken 22-VI-83, sampled 12-VII-83
9	MT-91-83	5564	26-IX-83	.8058	3.0	1.3	1.7	7.7	5.2	0.8	0.2	146	2.6	56.2	107	70	98	25	55.4	49.2	9	Taken in poundnet off Cherry point, in Chesapeake Bay. Apparently healthy, held at Lab prior to sampling.
16	MT-12-84	5752	29-V-84	.7970	1.6	0.3	1.3	3.2	5.9	1.1	0.1	149	3.0	49.6	110	71	104	65	78.8	64.4	3	Taken in Lynnhaven Inlet at the Bay Mouth, high salinity. Unhealthy, covered in Barnacles and emaciated, held at VIMS for treatment. Low hematocrit
22	MT-32-84	5761	6-IV-84	.7980	2.8	0.9	1.9	2.0	3.9	0.6	0.1	158	3.1	51	109	77	107	30	59.5	49.1	3	Taken from Lynnhaven Inlet, high salinity area, apparently healthy, slight heat stress, active
24	MT-62-84	5763	14-VII-84	.7966	2.9	1.3	1.6	4.8	5.3	1.0	0.3	164	3.6	45.5	116	78	113	53	71.8	58.4	7	Apparently healthy, from poundnets on York Spit some heat stress,
49	MT-124-84	5805	22-VII-85	.7985	4.1	1.9	2.2	8.7	5.9	0.6	0.1	164	3.8	43.2	113	79	111	32	63.5	50.4	10	Sick turtle from Deltaville, emaciated respiratory compromised. A later sample taken after holding this animal for 10 mos. 22-VIII-85
54	MT-137-84	5808	25-X-84	.7911	3.6	1.3	2.3	8.6	5.0	0.6	0.2	163	4.0	40.7	116	80	115	39	63.7	52.1	12	Apparently healthy, taken from Potomac River, cloacal temp. 21°C,

Table 12 . Intermediate Turtles Described by Factor Analysis

ID no.	Marine Turtle No.	Sam-ple No.	Date	Factor Score	Tpro (g/dl)	alb (g/dl)	Glob (g/dl)	Ca (mg/dl)	Phos (mg/dl)	Uric acid (mg/dl)	Cre (mg/dl)	Na (meq/l)	K (meq/l)	Na/K	Cl (meq/l)	BUN (mg/dl)	Glu (mg/dl)	Wt (kg)	CLS (cm)	CWS (cm)	Zone	Comments
60	MT-16-85	5859	16-V-85	.7999	3.9	1.2	2.7	7.7	3.4	0.7	0.1	157	2.9	54.1	107	78	105	52	73.7	58.5	12	Apparently healthy taken in the Potomac River, cloacal temp. 24°C
62	MT-19-85	5863	22-V-85	.7999	4.6	1.9	2.7	10.4	6.0	1.0	0.2	161	4.1	39.3	107	75	107	22	51.5	45.9	12	From Potomac River, fat apparently healthy, no epibiota.

with some input from sodium and urea nitrogen. Turtle MT-65-84 was recaptured three separate times over the summer of 1984. Its blood samples were included as separate samples. When these values were examined in the analysis they progressed from one group to another with each subsequent sample taken later in the season. Animals from endmember group one were characterized as coming from high salinity (>30 ppt) and having low cloacal temperatures (<22°C). Sera partial profiles for turtles with low body temperatures are found in Table 13. Animals found in endmember group 2 were generally (except MT-54-85) listed as stressed, sick or had abnormally high cloacal temperatures.

Table 13.  
Sera Profiles for *Caretta* with Low Body Temperatures  
( $<18^{\circ}\text{C}$ )

ID No.	Turtle Number	Date	total protein	albumin	calcium	phosphorus	sodium	potassium	chloride	urea nitrogen	glucose	Temp. $^{\circ}\text{C}$
5851	MT-02-85	09-V-85	2.9	1.1	8.2	5.5	164	4.2	118	27	72	16.0
5898	NC-CC	19-II-85	3.3	1.3	6.1	5.9	158	2.9	109	70	118	
5852	MT-03-85	09-V-85	3.3	1.0	8.5	4.3	164	4.2	116	31	84	18.3
5853	MT-04-85	09-V-85	4.4	1.4	10.3	5.2	164	3.6	116	26	76	17.0
5809	MT-140-84	13-XII-84	4.2	1.2	8.0	4.6	163	2.3	119	20	57	15.9 (8.2 $^{\circ}\text{C}$ on 12-XII-84)
5810	MT-140-84	20-XII-84	4.0	1.2	7.6	5.8	163	2.5	118	36	120	18.0

## V. DISCUSSION

### Underlying General Trends

Trends for serum values of turtles from Chesapeake Bay can be broken into several related categories. Serum measurements were affected by geographic location (due to physical differences in sites), seasonal changes, and pathological differences. Trends in serum samples primarily indicated effects on two major systems. The systems most reflecting changes were osmoregulation and metabolism (or nutritional differences). Results of a multi-variate analysis of the data also support these observations.

### Geographic Variations

Animals from the northern site were most distinctly different from animals from the southern site. These sites were the high salinity (30-32 ppt) southern Bay mouth site at Lynnhaven Inlet (Zn 3, Figure 1) and the more northern lower salinity (8-18 ppt) site at the Potomac River mouth (Zn 12) (Chen et al. 1984; Kuo et al. 1982). Sites located between these areas were more closely related to the southern site. Animals from southern sites were exposed to higher salinity and most samples were taken earlier in the season than samples from animals at the northern site. Because of geographic and physical differences in location, behavior of turtles differed between sites (Bellmund et al., 1987). Animals in the northern site were more representative of resident feeding populations that move into the Bay



early during the season and remain until cold weather forces migration out of the Bay (Lutcavage and Musick, 1985; Keinath et al , 1988). This behavior exposes these animals to longer periods of lower salinity.

Animals from the northern area were taken from nets that use fishing methods different than those of the southern Bay. These nets have a large by-catch of the Blue crab, Calinectes sapidus. Sea turtles taken from these nets had feces primarily composed of this species of crab. Caretta taken from more southern sections of the Bay are documented to be primarily feeding on the horseshoe crab, Limulus polyphemus (Lutcavage and Musick, 1985; Lutcavage, 1981). The choice of prey items may impose variable nutritional differences between locations.

#### Osmoregulation

Physiologically, sea turtles, as regulating poikilothermic animals, must compensate for a wide range of environmental changes. Ionic regulation affects acid-base balance, water regulation, and internal homeostatic control of many systems (Lutz, 1972; Stewart, 1978; Dunson and Weymouth, 1965; Dunson, 1979; Jackson and Ultsch, 1982). Data taken from the northern site show that animals from this area have lower but more variable electrolyte levels than animals taken from the higher salinity site (Table 4). This may reflect lower activity of the salt gland in response to decreased osmotic challenge. Evans (1973) found that sodium efflux from Caretta was 300uM/100g/hr in sea water. After transfer from sea water (SW) to



fresh water (FW) gland function decreased and sodium efflux decreased by 90%-99% within 6 hr and attained a rate of 0.7-1.4uM/100g/hr or 0.2-0.5% of the sodium loss of seawater acclimated animals after 1-2 days (Evans, 1973). Salt gland function in Chelonia mydas shut down after transfer from sea water to fresh water within 25 hours (Kooistra and Evans, 1976). Salinity in zone 12 ranges from 8 ppt to 18 ppt from April to Dec depending upon rainfall (Chen et al., 1984).

Animals foraging over the season in lower salinity can be expected to have lower electrolyte values and different sodium:potassium ratios than animals from higher salinity waters. Although salt gland function shuts down in freshwater to retain electrolytes, sodium efflux continues via the cloaca and integument (Kooistra and Evans, 1976; Robinson and Dunson, 1976; Evans, 1973). Grigg (1981) found that Crocodylus porosus collected from the the wild over a salinity gradient from fresh water to sea water had, in freshwater, decreased serum sodium, increased serum potassium, and decreased urinary sodium. In salt water these animals had increased serum sodium, decreased serum potassium, and increased urinary potassium. Turtles from the Potomac River (low salinity) showed lower serum sodium and chloride averages than the group for Virginia animals but their average potassium concentration increased over the summer. Turtles in Lynnhaven (Zone 3, the mouth of the Bay) showed higher and more constant levels of serum sodium and chloride but lower levels of potassium, reflecting the trends reported by Grigg (1981) in C. porosus. When serum values for all animals were grouped

together sodium and chloride declined with time over the summer. This decline probably reflects extended periods of time in lower salinity.

#### Nutritional Differences

The sodium changes in these animals may also reflect the decreased input of sodium through nutritional pathways (Dunson and Mazzotti, 1988). The cutaneous uptake of sodium by the diamond-backed terrapin (Malaclemys terrapin) was found to be negligible and the major route of sodium uptake in seawater occurred orally, through the buccal cavity (Robinson and Dunson, 1976). Evans (1973) noted that in the loggerhead reduced salt load was followed by a reduction in nasal salt gland secretions. The reduction in salt load acts as a feedback mechanism reducing efflux of salt from the system, however continued integumental and cloacal efflux occurs. If this occurs in Chesapeake Bay turtles then animals not actively taking in salt would still excrete it at some level. Although the salt gland ceases to pump ions, there is still ionic flux via the integument and the cloaca (Kooistra and Evans, 1976; Robinson and Dunson, 1976; Evans, 1973). In support of this theory, turtles from the Potomac River and hibernating turtles examined by Lutz and Dunbar-Cooper (1987) from the Florida coast also had lower sodium values than animals actively eating in high salinity waters.

Animals taken from the northern site had consistently higher values for serum total protein, albumin, urea nitrogen, and glucose than animals from the southern sites (Figures 6 & 8). In addition

these animals were generally larger and heavier with average  $Q$   $[(Q=CLS3/wt) \times 10^{*5}]$  values that were higher than animals from the southern sites (Bellmund et al., 1986). General observation of the pectoral and pelvic areas of these animals indicated that they were fatter and apparently healthier than animals from more southern sites. This may be affected by the difference in diet with the northern animals feeding more extensively on Callinectes while southern animals fed primarily on Limulus (Lutcavage, 1981; Bellmund et al., 1987).

#### Seasonal Variations

##### Osmoregulation

Seasonal affects are related to length of time animals are exposed to salinities less than that of full strength sea water. This is one reason for differences observed in apparently healthy animals from the northern Potomac River site and other southern sites examined. Explanations for these observed trends include an increase in fluid output. In fresh water cloacal fluid has a flow rate of 2-4 times that in SW (Kooistra and Evans, 1976). In low salinity water animals may increase their intake of water, which in conjunction with salt gland shut down, help maintain sodium and electrolyte levels.

Caretta in the Potomac River show an inverse relationship between sodium and potassium over time (Figure 7). Animals in lower salinity water continue to exchange sodium over an entire summer season even though the osmoregulatory challenge is severely reduced. The seasonal results of this adaptation are that animals remaining in

relatively high salinity waters (>18ppt) for the summer have higher serum electrolyte levels than animals foraging in lower salinity waters for the entire season. This is the pattern found during the summer season in Chesapeake Bay animals

This interpretation is supported by Bennett et al. (1986) finding that hatchling C. caretta drink sea water to compensate for osmotic and excretory water loss. In Crocodylus porosus plasma potassium was different (4.1-3.5mM/L) over a salinity gradient from fresh water to sea water (Grigg, 1981). In this same study potassium in cloacal urine of C. porosus showed a steady increase proportional to the salinity gradient. If C. caretta follow the same strategy as Malaclemys and C. porosus they then would excrete potassium, which is taken up as a counter ion to sodium, via the kidney (Robinson and Dunson, 1976; Gilles-Baillien, 1973; Grigg, 1981). If this were true then one would expect the sodium:potassium ratio to change over environmental salinity ranges. The Na:K ratio for these animals by location is found in Table 4. This ratio demonstrates compensatory regulation of sodium and potassium for C. caretta from Chesapeake Bay, in C. caretta studied by Evans (1973), and in Caretta from Florida (Lutz and Dunbar-Cooper, 1987)

#### Metabolism and Excretion

In addition to salinity differences, hibernation, decreased feeding, activity, and temperature also have an important affect on osmoregulation and blood chemistry of sea turtles. Lutz and Dunbar-Cooper (1987) have defined the biochemical plasma profile of an

apparently hibernating sea turtle. The most definitive of these parameters were believed to be sodium, magnesium, and calcium (Lutz and Dunbar-Cooper, 1987). Seasonal changes are also related to variations in temperature from early season low temperatures during migration to higher temperatures in late summer.

Changes seen by Haggag et al (1966) indicate that protein, glucose, and non-protein nitrogen change during hibernation in the lizard Varanus greseus. They found that there was a decrease in protein and glucose and an increase in non-protein nitrogen in hibernating animals. These trends mirror trends seen in early season animals in Chesapeake Bay and also reflect changes seen by Lutz and Dunbar-Cooper (1987). Animals in Chesapeake Bay were lower in total protein, inorganic phosphorus, and urea nitrogen early in the season. Haggag et al. (1966) attribute the rise in non-protein nitrogen to a build up of excretory end products due to decreased excretion during hibernation. Animals in Chesapeake Bay forage for the summer season and would be expected to increase all of these parameters to some extent in response to food availability and seasonal temperature. Temperature affects digestive turnover rate and food intake in reptiles and so an animals digestive patterns and efficiency are directly affected by seasonal temperature fluctuations (Parmenter, 1981). Animals migrating into Chesapeake Bay early in the season have moved through water that is from 14-16oC (Musick unpubl observation). This means that animals moving into the Bay early in the season are physiologically acclimatized to temperatures that are much lower than the temperature of the water they were taken from.

Turtles would then be expected to exhibit serum profiles more characteristic of cold reptiles, that is animals with a lowered metabolic rate. Figure 14 shows the results of work on one Caretta which was sampled after feeding to find the affects of lowered temperature on glucose and urea nitrogen metabolism. This work showed that decreased temperature changed relative circulating levels of urea nitrogen and glucose after feeding. This same relationship was also visible in animals arriving early in the season in Chesapeake Bay and in cold stunned animals from this region. Cold stunned animals have low urea nitrogen and glucose values relative to mean values of these parameters calculated for apparently healthy animals (Tables 7 & 14). Jackson and Ultsch (1982) found that at 30°C potassium increased from 2.4 meq/l to 9.0 meq/l in anoxic freshwater turtles (Chrysemys picta belii). They also found dramatic increase in total calcium and magnesium.

Gilles-Baillien (1973) reports that Malaclemys acclimatized to seawater had higher plasma osmotic pressure than animals acclimatized to fresh water. This increase in osmotic pressure is attributed to an increase in sodium, chloride and urea in sea water adapted animals (Gilles-Baillien, 1970). Urea values were found to peak in September, just prior to hibernation, in seawater adapted Malaclemys and this effect was described as an osmoregulatory adaptation to compensate for hibernation decreases in other electrolytes (Gilles-Baillien, 1973). This pattern was also seen in loggerhead sea turtles from Chesapeake Bay whose glucose and urea nitrogen increased during the summer until September. The patterns in Malaclemys for



sodium, chloride, and potassium were similar for both fresh and salt water adapted animals, these values were highest from May to September but were progressively lowered during hibernation (Gilles-Baillien, 1973). Loggerheads in Chesapeake Bay showed an increase over the summer in total protein, albumin, and phosphorus.

#### Pathological Changes

Sick and injured animals had a lower mean sodium level than apparently healthy animals. If sodium, chloride, and potassium are plotted together in apparently healthy animals and in sick animals, sodium and chloride change together in healthy animals but all three electrolytes change together in sick or injured animals. Sick and injured animals generally had serum components that were more variable for electrolytes and total protein/albumin values. This group also had higher mean globulins and lower mean urea nitrogen and glucose than apparently healthy animals.

Pathologic changes in these animals can be related to temperature effects, disease affects, and changes related to severe emaciation (Sturbaum and Bergman, 1981; Austin et al., 1927; Bonnet, 1979; Frye, 1981). Temperature changes affect the rate of uptake and excretion involved in renal processing, and digestive or metabolic function. Disease affects vary from animal to animal. Severely emaciated animals often had high serum total protein and low or extremely variable electrolyte values. Emaciation may, simply be a manifestation of internal disease effects and result in high globulin values and severely altered osmoregulatory capacity. Lutz et al

(1986) found that the main system compromised by application of crude oil to sea turtles was osmoregulation. However complex physiological interrelationships function to control metabolic processes and are often not detectable by examination of individual mean values in blood of lower vertebrates (Lutz,1972)

### Multi-Variate Analyses

#### Principal Component Analysis

Principal component analysis (PCA) of the data was run on animals classified as apparently healthy or sick and injured. Three factors groups were defined by PCA for both groups of animals. The first factor in both groups was composed of the variables albumin, total protein, calcium, and phosphorus. The second factor group in healthy animals was composed of urea nitrogen, potassium, and glucose. Factor group 3 was composed of sodium and chloride. PCA of sick and injured animals defined factor groups for factors two and three differently than they were defined for healthy animals. Group 2 was composed of sodium, chloride and potassium. Group 3 was composed of urea nitrogen and glucose.

The first factor in PCA was defined by the same variables in both healthy and sick groups. This result indicates that this is a fundamentally important group and is stable whether the animals are healthy or not. Calcium and phosphorus interact with all ions and charged species to affect ionic regulation and acid-base balance (Stewart, 1978; Jackson and Ultsch, 1982). These compounds can be present in blood, in part bound to proteins, particularly albumin.



They are affected in juvenile animals by variations in ionic regulation, acid base balance, growth requirements, temperature, metabolism, and various pathologies (Guyton, 1982; Frye, 1981; Sturbaum, 1982).

Since all animals in this work were juvenile, growth would also be expected to be an important component of serum chemistry. Growth regulates the circulating concentrations of calcium, and phosphorus. As juvenile animals grow bone is laid down at a rapid rate. In chelonians the massive amount of bone laid down makes the circulating concentration of calcium very important in the total changes in circulating biochemical parameters examined. Calcium is carried by plasma proteins such as albumin. The calcium/phosphorus ratio in bone is between 1.3 and 2.0 (Guyton, 1982). The circulating and incorporated concentration of calcium is regulated by a variety of mechanisms and is dependent on phosphate as a balancing compound. Hence, as expected, calcium and phosphorus in Chesapeake Bay animals were related to total protein and albumin in PCA. In addition the ontogenic changes seen in the shells of developing juvenile chelonians is responsible for changing availability in the exchangeable ion pool (Dunson and Heatwole, 1986). This is important for calcium and phosphorus regulation as well as for sodium and potassium.

Changes in serum concentrations of calcium or phosphorus may come from disruption of internal regulatory mechanisms, physiological changes due to a varying physical environment (ie. temperature and

salinity), or from nutritional input of these compounds. The presence and circulating concentration of ionized calcium is determined by the nutritional availability of calcium, the availability and activity of vitamin D3 (cholecalciferol), temperature, the ionic balance of the plasma, the colloidal osmotic pressure from calcium/albumin complexing, and the absorption or removal of calcium by the body (Frye, 1981; Jackson and Ultsch, 1982; Guyton, 1981; Jackson et al , 1984; Stewart, 1978). Many of these conditions combine to result in synergistic or concurrent effects and can often be interrelated. Calcium in reptiles is involved in bone growth, membrane function, and the strong ion difference (SID) (Jackson et al , 1984; Stewart, 1981; Ultsch et al , 1984). Calcium is used as part of the buffering system within turtles (Jackson et al , 1984; Ultsch et al , 1984) and can be mobilized from bone.

Changes in the composition of variables in the second factor group in both data sets (for apparently healthy and sick or injured turtles) were important in examining internal homeostatic changes and differences between these two groups. In apparently healthy animals the second Factor group was composed of urea nitrogen, potassium, and glucose. In sick and injured animals, the second Factor group was composed of sodium, chloride, and potassium. The composition of Factor group 2 for apparently healthy animals was made up of parameters that are controlled primarily by excretion and metabolism. Factor group 3, for apparently healthy animals, was composed of sodium and chloride, a group obviously related to osmoregulation.

Factor group 2 on sick and injured animals was made up of sodium, chloride, and potassium. Factor group 3 was made up of urea nitrogen and glucose. The switch of importance of potassium in this group of animals from the group which apparently describes renal activity and includes urea nitrogen and glucose to the osmoregulatory group with sodium and chloride indicates an important internal change. The shift in these variable groupings indicates that osmoregulatory ability was affected in the sick group. The change in electrolyte relationships could be due to a loss of function in the salt gland and the loss of the Na-K pump. The same effect was seen in the application of crude oil to sea turtles, this resulted in the loss of osmoregulatory capacity (Lutz *et al.*, 1986). Due to these effects it appears that the most important system to examine in sea turtles to determine the overall status of an animal is to examine the electrolyte balance and the Na<sup>+</sup>, Cl<sup>-</sup>, and K<sup>+</sup> concentrations

#### Q-Mode Analysis

Turtle groupings in Q-mode analysis were based primarily on salinity, temperature, and physiological stress. The variables delineating these groups were the same. In both cases sodium, urea nitrogen, and glucose (chloride may also be included, depending upon cut-off values for Varimax factor loadings) were included. Osmoregulation and temperature were of primary importance. Gilles-Baillien (1973a&b) and Haggag *et al.* (1965, 1966), among others, report the use of non-protein nitrogen in reptiles as an osmoeffector. In addition, they report that all components picked in this analysis change with hibernation and seasonality. Increased

glucose in the endmember group for Factor II can be considered as a response to heat stress or disease state and the resulting physiological changes that occur in moribund or stressed animals (Sturbaum, 1982; Sturbaum and Bergman, 1981; Frye, 1981; Paulson and Hutchinson, 1987). Wedemeyer (1970) describes Selye's general adaptive syndrome in response to stress as including increases in blood glucose level and nitrogen metabolism. The general adaptive syndrome may be characterized by non-specific metabolic changes such as " The morphological, biochemical, and physiological changes which occur as a result of stress " (Wedemeyer, 1970). Changes accompanying this syndrome include increased nitrogen metabolism and increased blood glucose level. Physical response is manifested by alarm reaction, resistance and exhaustion. These changes could result in the grouping of apparently stressed animals with elevated internal temperatures and high serum glucose into Factor group II.

Three dying animals were found in the Factor group II list of end members. Turtle MT-113-83 died shortly (within a week) after blood was drawn for analysis. This animal was severely emaciated yet had the second highest blood glucose level of all animals examined. All sick turtles in this factor group had low blood urea nitrogen levels. Stress hormones in these animals were not assessed due to the uncertainty of the results and because 1983 analyses were not available. Animals included in Factor group II due to increased temperatures were generally characterized by fighting and  $>30^{\circ}\text{C}$  temperatures. It is possible that animals were included due to

nutritional status but it was not possible to determine the magnitude of this effect

Animals from Factor group I were characterized by low cloacal temperatures ( $<22^{\circ}\text{C}$ ) and being found in high salinity (generally off-shore) waters. This group of animals included two severely cold stunned animals ( $< 10^{\circ}\text{C}$ ) that plotted slightly off the factor line. All animals, except those that were cold stunned, had high urea nitrogen and low glucose. No obvious trend was found for absolute values of sodium and chloride, however they were included as the two most important variables in this Factor group. When serial urea nitrogen and glucose measurements were made on a single animal held at different temperatures their relationships changed. At  $20^{\circ}\text{C}$ , urea nitrogen was higher than glucose but at  $26^{\circ}\text{C}$  this relationship was reversed. These changes in these components with temperature nicely reflect relationships seen in Factor group I animals from the field.

Factor scores were tightly grouped for a third set of animals. Factor scores in this group ranged from .7900-.8100 . This group of animals include two sub-clinically ill turtles and other animals that had no obvious reason to be included. It may be that this group represents the sub-clinically ill group of animals that have no other obvious symptoms other than complex internal changes.

## VI. CONCLUSIONS

This work provides a data base of blood serum values for healthy and sick loggerhead sea turtles from the wild. It illustrated the difficult nature of assessing condition from absolute serum values in wild populations with no previous history. It was not possible to divide turtles into sick and healthy categories based upon individual serum profiles. Indeed, it was not possible to divide turtles into these categories using multivariate analyses. Using multivariate analyses it was only possible to look at relative relationships within serum variables and develop categories based upon variable relationships. Profile groups using PCA were based on metabolism and growth, ionic environment, and metabolic excretory state. Factor analysis divided groups based upon their physical environment and in part on their health (internal environment). It showed the importance of external physical environment to poikilothermic animals and the inherent difficulties in describing animals closely tied to their external environment.

Relationships and general relative differences in various serum compositions were reflected in all analyses and the literature. Multivariate factor analysis divided turtles along a gradient best defined by salinity, temperature, and physiological stress. Data and

analyses suggest that osmoregulation is an important system to examine when evaluating internal homeostasis in marine turtles. Marine animals must have an effective functional osmoregulatory system to continue to maintain a favorable osmotic gradient. Once this system is compromised the entire well being of the animal is in jeopardy. As such this system is both conservative in its physiological range and responsive to environmental fluctuations in temperature, nutritional availability, and particularly salinity. The result of this dual nature is a system that is critical in maintaining internal homeostasis and valuable in determining the status of the animal.



Appendix I.  
Serum Analytical Methods

Test Name	Catalog Number	Application Sheet Reference Number
Total Protein	749117	6-417-0285
Albumin	620174	8-50414501-1282
Calcium,GABA	857825	6-689-0887
Phosphorus	836281	6-443-0186
BUN/Glucose Twin Reagent	842273	6-951-0787
Uric Acid	704156	6-477-0684
Creatinine	704130	6-494-0684
Total Bilirubin, DPD	977179	6-495-0687
GOT/GPT Twin Pack	842265	6-592-0486
Alkaline Phosphatase	704024	6-455-0587
LDH, TRIS	620099	8-50415001-1282
Na, K, Cl, by Ion Selective Electrode	749044, 749052, 836105, 836113, 836121, 646911, 646938, 608250	6-457-0187

Reference Book Title: Boehringer Mannheim  
Diagnostics, Inc. Hitachi 705, Operator's Reference  
Manual, Section 3. REAGENTS

Published by ;  
Boehringer Mannheim Diagnostics, Inc.  
Hitachi 705 Clinical Support  
9115 Hague Road  
Indianapolis, Indiana 46250



Appendix II

Test	Method	Reference
Blood Urea Nitrogen (BUN)	Quantitative colormetric determination of BUN - Direct method	Crocker, C.L., A. J. Med. Tech., 33,361 (1967)
Glucose	Quantitative colormetric determination of glucose.	Hultman, E. Nature 183,108. 1959. Dubowski, K.M. Clinical Chem. 8, 215,1962.

## VII. LITERATURE CITED

- Abel, J. H. and R. A. Ellis. 1966. Histochemical and electron microscopic observations on the salt secreting lacrymal glands of marine turtle. *Am. J. Anat.*, 118, 337-358.
- Austin, J. H., F. W. Sunderman, and J. G. Camack. 1927. Studies in serum electrolytes: II. The electrolyte composition and the pH of serum of a poikilothermous animal at different temperatures. *J. Biol. Chem.* 72:677-685
- Bacon, P. R. 1981. The status of sea turtle stocks management in the Western Central Atlantic. Western Central Atlantic Fishery Commission, UNDP, WECAF Studies No. 7. 38pp.
- Bellmund, S. A., J. A. Musick, R.E. C. Klinger, R. A. Byles, J. A. Keinath, and D. E. Barnard. 1987. Ecology of Sea Turtles in Virginia, Special Scientific Report No. 119, Va. Inst. of Marine Sci., Coll. William and Mary, Gloucester Pt., Va 23062.48 p.
- Bennett, J. M. , L. E. Taplin, and G. C. Grigg. 1986. Sea water drinking as a homeostatic response to dehydration in hatchling loggerhead sea turtles Caretta caretta. *Comp. Biochem. Physiol.* 83A, (3), 507-513.
- Bentley, T. B. and A. Dunbar-Cooper. 1980. A blood sampling technique for sea turtles. Contract No. Na-80-GE-A-00082 for National Marine Fisheries Service, September 1980.
- Berquist, C. R. 1986. Stratigraphy and Heavy Minerals in the Lower Chesapeake Bay. Unpublished Dissertation from Va. Inst. Marine Sci., Coll. William and Mary. 105 p.
- Bonnet, B. 1979. Influence of the nutritional conditions on the organic composition of blood and urine in the juvenile sea turtle Chelonia mydas L. *Aquaculture*, 16, 253-260.
- Brock, J. A., R. M. Nakamura, A. Y. Miyahara, and E. M. L. Chang. 1976. Tuberculosis in Pacific green sea turtles Chelonia mydas L. *Trans. Am. Fish. Soc.*, 105 (4), 564-566.
- Byles, R. A. In Prep. Diving Behavior in Virginia Sea turtles Caretta caretta and Lepidochelys kempi.
- Carr, A. 1952. Handbook of Turtles. Comstock publishing associate Cornell University press, London, 535 p.

- Carr, A., L. Ogren, and C. McVea. 1980. Apparent hibernation by the Atlantic loggerhead turtle (Caretta caretta) off Cape Canaveral, Florida Biol. Conserv. 19; 7-14.
- Chen, H. S., P. V. Hyer, Y. Unkulvasapaul, and B. Neilson. 1984. Summary Report on the Calibration of the Water Quality Models of the Chesapeake Bay System. VA. Inst. of Marine Science, Gloucester Pt. Va., College of William and Mary. pp. 79.
- Coulson R. A. and T. Hernandez. 1980. Alligator Metabolism; Studies on Chemical Reactions in vivo. Pergamagon Press, London.
- Davis, J. C. 1986. Statistics and Data Analysis in Geology. 2ND ED. John Wiley and Sons, New York, 646pp.
- Dessauer, H. C. 1970. Blood chemistry of reptiles: physiological and evolutionary aspects . In: C. Gans and T. S. Parsons (eds) Biology of the Reptilia. p. 1-72, Morphology 3, Acad. Press, N.Y.
- Dunson, W. A. 1967. Sodium fluxes in freshwater turtles. J. exp. Zool., 165:171-182.
- Dunson, W. A. 1970. Some aspects of electrolyte and water balance in three estuarine reptiles, the diamondback terrapin, American and "salt water" crocodiles. Comp. Biochem. Physiol. 32:161-174.
- Dunson, W. A. 1976. Salt glands in reptiles, In: Vol. 5. C. Gans and W.R. Dawson (eds.) Biol. of the Reptilia, Academic Press, N.Y. pp. 413-445.
- Dunson, W. A. 1978a. Role of the skin in sodium and water exchange of ( aquatic snakes placed in seawater. Am. J. Physiol. 235 3):151-159.
- Dunson, W. A. 1978b. Salt and water balance in sea snakes, In: Biology of the Sea Snakes. W. A. Dunson (ed.)
- Dunson, W. A. 1979. Control mechanisms in reptiles In: Mechanisms of Osmoregulation in Animals, R. Gilles (ed.) Wiley Interscience, N.Y., pp. 273-322.
- Dunson, W. A. 1984. Contrasting roles of the saltglands, the integument, and behavior in osmoregulation of marine and estuarine reptiles. In: Osmoregulation Symposium, Fifth Conf. European Soc. Comp. Physiol. Biochem., R. Gilles and A. Pequeax eds., Springer.
- Dunson, W. A. 1985. Effect of water salinity and food salt content on growth and sodium efflux of hatchling diamondback terrapins (Malaclemys). Physiol. Zool. 58(6):736-747.

- Dunson, W. A. 1986. Estuarine populations of the snapping turtle Chelydra as a model for the evaluation of marine adaptations in Reptiles. *Copeia* 1986 (3):741-756.
- Dunson, W. A. and J. Freda. 1985. Water permeability of the skin of the amphibious snake. Agkistrodon piscivorus. *J. of Herpetology* 19 (1):93-98.
- Dunson, W. A. and H. Heatwole. 1986. Effect of relative shell size in turtles on water and electrolyte composition. *Rapid Communications. Am. J. Physiol.* 250 (Regulatory Integrative Comp. Physiol. 19).
- Dunson, W. A. and F. J. Mazzotti. In press. Salinity as a limiting factor in the distribution of reptiles in Florida Bay: A theory for the estuarine origin of marine snakes and turtles. *Bull. Mar. Sci.*
- Dunson, W. A. and E. O. Moll, 1980. Osmoregulation in sea water of hatchling Emydid turtles, Callagur borneoensis, from a Malaysian sea beach. *J. Herp.* 14(1):31-36.
- Dunson, W. A. and R. D. Weymouth. 1965. Active uptake of sodium by softshelled turtles (Trionyx spinifer), *Science*, 149:67-69.
- Ellis, R. A. and J. H. Abel Jr. 1964. Intercellular channels in the salt-secreting glands of marine turtles. *Science* 144:1340-1343.
- Ellis, T. M. 1981. Tolerance of sea water by the American crocodile Crocodylus acutus, *J. Herp.* 15:187-192.
- Evans, D. H. 1973. Sodium balance of the euryhaline marine loggerhead turtle Caretta caretta. *J. comp. Physiol.* 83, 179-185.
- Frair, W. 1977b. Sea turtle blood cell parameters correlated with carapace lengths. *Comp. Biochem. Physiol.*, 56, 467-472.
- Franzmann, A. W. and C. C. Schwartz. 1988. Evaluating condition of Alaskan black bears with blood profiles. *J. Wildl. Manage.* 52(1):63-70.
- Frye, F. L., 1981. Biomedical and Surgical Aspects of Captive Reptile Husbandry. *Veterinary Medical Publishing Company*, Edwardsville, Kansas. pp. 456.
- Full, W. E., Ehrlich, R. and Klován, J. E. 1981. EXTENDED QMODEL - Objective definition of external end-members in the analysis of mixtures. *J. of Math. Geology*. 13(4):331-344.
- Gilles-Baillien, M. 1970. Urea and osmoregulation in the diamondbacked terrapin Malaclemys centrata centrata (Latreille). *J. Exp. Biol.* 52, 691-697.

- Gilles-Baillien, M. 1973a. Isosmotic regulation in various tissues of the diamondbacked terrapin Malaclemys centrata centrata (Latreille). J. Exp. Biol. 59, 39-43.
- Gilles-Baillien, M. 1973b. Hibernation and osmoregulation in the diamondbacked terrapin Malaclemys centrata centrata (Latreille). J. Exp. Biol. 59, 45-51.
- Gilles-Baillien, M. 1973c. Seasonal variation in weight and hematological data of Testudo hermanni hermanni Gmelin. Arch. Internat. Physiol. Bioch., 81, (4), 723-732.
- Gilles-Baillien, M. 1974. Seasonal variation in reptiles. In: M. Florkin and B. T. Scheer (eds) Chemical Zool. IX:353-376. Acad. Press. N.Y.
- Grigg, G. C. 1981. Plasma homeostasis and cloacal urine composition in Crocodylus porosus caught along a salinity gradient. J. Comp. Physiol. 144B, 261-270.
- Grigg, G. C., L. E. Taplin, P. Harlow, and J. Wright. 1980. Survival and growth of hatchling Crocodylus porosus in salt water without access to fresh drinking water. Oecologia. 47:264-266.
- Gould, S. J. 1982. The Mismeasure of Man. Johns Hopkins Press. Baltimore, Md. pp. 352.
- Haggag, G., K. A. Raheem and R. Kahlil, 1965. Hibernation in reptiles. I. changes in blood electrolytes. Comp. Bioch. Physiol. 16:427-465.
- Haggag, G., K. A. Raheem, and F. Khalil. 1966. Hibernation in reptiles II. Changes in blood glucose, haemoglobin, red blood cell count, protein and non-protein nitrogen. Comp. Biochem. Physiol. 17:335-339.
- Haggag, G., K.A. Raheem and F. Khalil. 1966. Hibernation in Reptiles. III. Tissue analysis for glycogen and high energy phosphate compounds. Comp. Bioch. Physiol. 17:341-347.
- Haines, H. and W. C. Kleese. 1977. Effect of water temperature on a herpesvirus infection of sea turtles. Infect. and Immun., 15 (3);756-759.
- Haines, H. 1978. A herpesvirus disease of green sea turtles in aquaculture, Mar. Fish. Rev., 40 (3);330-37.
- Hildebrand, H. H. 1981. A historical review of the status of sea turtle populations in the western Gulf of Mexico. In: K. A. Bjorndal ed., Biology and Conservation of Sea Turtles. Proceedings of the World Conference on Sea Turtle Conservation, Washington D. C., 26-30 November 1979. Smithsonian Institution Press, Wash. D.C., pp. 447-453.

- Hillestad, H.O., J.I. Richardson, C. McVea, Jr., J. M. Watson, Jr.  
1981. Worldwide Incidental capture of sea turtles. In: K. A. Bjorndal ed., *Biology and Conservation of Sea Turtles*. Proceedings of the World Conference on Sea Turtle Conservation, Washington D. C., 26-30 November 1979. Smithsonian Institution Press, Wash. D.C., pp. 489-496.
- Holmes, W. N. and R. L. McBean. 1964. Some aspects of electrolyte excretion in the green turtle Chelonia mydas L. *J. Exp. Biol.* 41:81-90.
- Hull, C. H., N. H. Nie. 1981. SPSS Update 7-9; New Procedures and Facilities for Releases 7-9. McGraw-Hill, N.Y., N.Y. pp. 402.
- Hutton, K. E. and C. J. Goodnight. 1957. Variations in the blood chemistry of turtles under active and hibernating conditions. *Physiol. Zool.* 30:198-207.
- Imbrie, J. and E. G. Purdy, 1962. Classification of modern Bahamian carbonate sediments. In: *Classification of Carbonate Rocks A Symposium: Amer. Assoc. Petroleum Geologists Mem.* 1:253-272.
- IUCN. 1971. Marine turtles, proceedings of the second working meeting of marine turtle specialists. Organized by the survival service commission, IUCN, 8-10 march 1971, Morges, Switzerland. IUCN publications new series, supplementary Paper no.31, 108 p.
- Jackson, D. C. 1985. Respiration and respiratory control in the Green turtle, Chelonia mydas. *Copeia* 3:664-671.
- Jackson, D. C., and G. R. Ultsch. 1982. Long-term submergence at 30°C of the turtle Chrysemys picta bellii in normoxic and severely hypoxic water. II. Extracellular Ionic responses to extreme lactic acidosis *J. Exp. Biol.* 96, 29-43.
- Jackson, D. C., C. V. Herbert, and G. R. Ultsch. 1984. The comparative physiology of diving in North American freshwater turtles. II. Plasma ion balance during prolonged anoxia., *Physiol. Zool.* 57 (6), 632-640.
- Jacobson, E. R., personal communication 26 February 1987, Seventh Annual Workshop on Sea Turtle Conservation and Biology.
- Joreskog, K. G., J. E. Klován, and R. A. Reymont. 1976. *Geological Factor Analysis*. Elsevier, New York, 178pp.
- Keinath, J. A., J. A. Musick, and R. A. Byles. 1987. Aspects of the biology of Virginia sea turtles: 1979-1986. *VA. J. Science*, 38(4):329-336.
- Klován, J. E. 1975. R and Q-Mode factor analysis, In; McCammon, R. B. ED. *Concepts in Geostatistics*, Springer Verlag, New York. (21-70p) 168pp.



- Klovan, J. E. and A. T. Miesch. 1975. Extended CABFAC and QMODEL computer programs for Q-Mode factor analysis and calculation of factor scores. J. of the International Assoc. for Math. Geol., 3(1):61-77.
- Kooistra, T. A. and D. H. Evans. 1976. Sodium balance in the green turtle Chelonia mydas L. in seawater and freshwater. J. Comp. Physiol., 107, 228-240.
- Kuo, A. Y., P. V. Hyer, and B. J. Neilson. 1982. A Mathematical Model for the Study of Stormwater Impacts on the Water Quality of Lynnhaven Bay, Virginia. VIMS Special Report no. 262 in Applied Marine Science and Engineering, pp. 146.
- Lutcavage, M. and J. A. Musick, 1985. Aspects of the biology of sea turtles in Virginia. Copeia. 2:449-456.
- Lutcavage, M., P.L. Lutz, and H. Baier. 1987. Gas exchange in the loggerhead sea turtle, Caretta caretta. J. Exp. Biol. 131:365-372.
- Lutcavage, M. 1981. Status of Marine Turtles in Chesapeake Bay and Virginia Coastal Waters. A thesis presented to the college of William and Mary, School of Marine Science. Virginia Institute of Marine Science. Gloucester Point Virginia.
- Lutz, P. L. 1972. Ionic patterns in the teleost. Comp. Biochem. Physiol., 42A:719-733.
- Lutz, P. L. 1975. Adaptive and evolutionary aspects of the ionic content of fishes. Copeia. 2:369-373.
- Lutz, P. L. 1985. The regulatory biology of sea turtles. Copeia. 3:662-663.
- Lutz, P. L., and T. B. Bentley, 1985. Respiratory physiology of diving in the sea turtle. Copeia. 3:671-679.
- Lutz, P. L., and A. Dunbar-Cooper, 1987. Variations in the blood chemistry of the loggerhead sea turtle, Caretta caretta. Fishery Bulletin 85:1 pp.37-43.
- Lutz, P. L., M. Lutcavage, D. Hudson and G. Bossert. 1987. Effects of exposure to oil, from; Final Report on Study of the Effects of Oil on Marine Turtles. Vol 2 Technical Report. Prepared for the U.S. Dept. Interior, MMS, MMS contract Number 14-12-0001-30063. pp. 93-148.
- Maginniss, L. A., S. S. Tapper, and L. S. Miller. 1983. Effect of chronic cold and submergence of blood oxygen transport in the turtle, Chrysemys picta, Respiration Physiology. 53:15-29.
- Martin, E. Personal Communication, phone conversation 17 Nov. 1988.2

- Masat, R. J. and X. J. Musacchia. 1965. Serum protein concentration changes in the turtle, Chrysemys picta. Comp. Bioch. Physiol. 16:215-225.
- Milstein, A. 1986. A multivariate approach to environmental zooplankton relationships in Maldonado Bay (Uruguay). Bolm. Inst. Oceanogr., S. Paulo, 34(unico):13-21.
- Morris, Y. A., 1982. Steroid Dynamics in immature sea turtles. MS Thesis. Texas A&M University.
- Morris, Y. A. and D. W. Owens. 1982. Corticosterone and stress in sea turtles. Amer. Zool. 22:956.
- Musick, J. A., R. Byles, R. Klinger, and S. Bellmund. 1984. Mortality and Behavior of Sea Turtles in the Chesapeake Bay. Summary report for 1979-1983. Contract NA80FAC00004. with National Marine Fisheries Service, NE Region. by the Virginia Inst. of Marine Science., Gloucester Pt. , Va. 52 p.
- Nie, N. H., C. H. Hull, J. G. Jenkins, K. Steinbrenner, and D. H. Bent. 1975. Statistical Package for the Social Sciences, Second Ed. McGraw Hill Book Co., N.Y., N.Y. pp. 675.
- Ogren, L. and C. McVea, Jr. 1982. Apparent hibernation by sea turtles in North American waters. In: K. A. Bjorndal (ed.) Biology and Conservation of Sea Turtles. Proc. World Conference on Sea Turtle Conservation. 26-30, Nov. 1979, Wash. D. C. Smithsonian Inst. Press, 583 p.
- Owens, D. W. 1978. A technique for determining sex of immature Chelonia mydas L. using a radioimmunoassay. Herpetologica, 34 (4), 986-998.
- Owens, D. W. and G. J. Ruiz. 1980. Obtaining blood and cerebrospinal fluid from marine turtles. Herpetologica. 36:17-20.
- Owens, D. W. and Y. A. Morris. 1985. The comparative endocrinology of sea turtles. Copeia. 3:723-735..
- Prange, H. D., 1985. Renal and extra-renal mechanisms of salt and water regulation of sea turtles: a speculative review. Copeia. 3:771-776.
- Prange, H. D. and D. C. Jackson. 1976. Ventilation, gas exchange and metabolic scaling of a sea turtle. Respiration Physiology, 27:369- 377.
- Paulson, B. K. and V. H., Hutchinson. 1987. Blood changes in Bufo cognatus following acute heat stress, Comp. Bioch. Physiol., 87A(2):461-466.



- Richardson, J. I., T. H. Richardson, and M. W. Dix. 1978. Population estimates for nesting female loggerhead sea turtles (Caretta caretta) in the St. Andrew Sound area of southeastern Georgia, U.S.A. Pages 34-38 IN G. E. Henderson, ed., Proceedings of the Florida and inter-regional conference on sea turtles, 24-25 July 1976, Jensen Beach, Florida. Fla. Mar. Res. Publ. No. 33. 66 pp.
- Robinson, G. D. and W. A. Dunson. 1976. Water and sodium balance in the estuarine diamondback terrapin Malaclemys, J. Comp. Physiol., B, 105:129-152.
- Ross, J.P. 1981. Historical Decline of Loggerhead, Ridley, and Leatherback sea turtles. In: K. A. Bjorndal ed., Biology and Conservation of Sea Turtles. Proceedings of the World Conference on Sea Turtle Conservation, Washington D. C., 26-30 November 1979. Smithsonian Institution Press, Wash. D.C., pp. 189-196.
- Schmidt-Nielsen, K. and R. Fange. 1958. Salt glands in marine reptiles. Nature, 4638:783-785.
- Schmidt-Nielsen, B., and L. E. Davis. 1968. Fluid transport and tubular intracellular spaces in reptilian kidneys. Science. 159:1105-1108.
- Schoffeniels, E. and R. R. Tercafs. 1965. Adaptation D'un retile marin. Caretta caretta L. al'eau douce et d'un reptile d'eau douce, Clemmys leprosa L. a l'eau der mer. Ann. Soc. R. Zool. Belg. 96:1-8.
- Seidel, M. E. 1975. Osmoregulation in the turtle Trionyx spiniferus from brackish and freshwater. Copeia. 1:124-128.
- Smith, H. W., 1929. The inorganic composition of the body fluids of the Chelonia, J. Biol. Chem., 82:651-661.
- SPSS. 1975 Statistical Package for the Social Sciences, 2 ND edition, C. H. Hull, G. G. Jenkins, K. Steinbrenner, and D. H. Bent. editors, McGraw Hill Book Co.
- Standora, E. A., J. R. Spotila, and R. E. Foley. 1982. Regional endothermy in the sea turtle, Chelonia mydas, J. Therm. Biol. 7:159-165.
- Stevens, R. personal communication 22 January 1987. Eastern States Veterenary Assciation Meetings, Orlando Florida.
- Sturbaum, B. A. 1982. Temperature regulation in turtles. Comp. Biochem. Physiol. 72(A), 4: 615-620.

- Sturbaum, B. A. and J. L. Bergman. (1981) Changes in selected blood components of the three-toed box turtle, Terrapene carolina triunguis, during heat stress. Comp. Bioch. Physiol. 70(A):599-602.
- Sturbaum, B. A. and M. L. Riedesel. 1974. Temperature regulation responses of ornate box turtles, Terrapene ornata, to heat. Comp. Bioch. Physiol. 48(A):527-538.
- Ultsch, G. R. and D. C. Jackson. 1982a. Long-term submergence at 30°C of the turtle Chrysemys picta bellii in normoxic and severely hypoxic water. I. Survival, gas exchange and acid base status. J. Exp. Biol. 96:11-28.
- Ultsch, G. R. and D. C. Jackson. 1982b. Long-term submergence at 30°C of the turtle Chrysemys picta bellii in normoxic and severely hypoxic water. III. Effects of changes in ambient Po<sub>2</sub> and subsequent air breathing J. Exp. Biol. 97, 87-99.
- Ultsch, G. R., C. V. Herbert, and D. C. Jackson. 1984. The Comparative physiology of diving in North American freshwater turtles. I. Submergence tolerance, gas exchange, and acid-base balance. Physiol. Zool. 57 (6), 620-631.
- Wedemeyer, G. 1970. The role of stress in disease resistance of fishes, pp. 30-35, In: A Symposium of Diseases of Fishes and Shell Fishes. Eds S. F. Snieszko. Special publication No. 5, American Fisheries Society, Washington, D. C.

## VITA

Sarah Anne Bellmund

Born in Washington, D. C. February 24, 1960. Graduated from Wakefield Senior High School in Arlington Virginia in 1978. Received B. S. in Biochemistry with a minor in Chemistry from the School of Arts and Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia in 1982. Entered the School of Marine Science, at the Virginia Institute of Marine Science, College of William and Mary in 1982. Leave of absence from school during the period 1985-1987 to work at the University of Maryland, Horn Point Laboratories as a Research Assistant.